

REGIONE DEL VENETO



Servizio Sanitario Nazionale – Regione Veneto
**AZIENDA UNITA LOCALE SOCIO-
SANITARIA N°3**
DIPARTIMENTO DI SCIENZE MEDICHE

U.O.C. Oncologia ed Ematologia Oncologica
(Direttore Dott. Giuseppe Azzarello)

Certificazione ISO 9001-2015
N 211181

LISTA DELLE PUBBLICAZIONI SCIENTIFICHE INTERNAZIONALI “PEER-REVIEWED”

Per ogni pubblicazione è riportato titolo e il nome della rivista scientifica con identificazione dei parametri di pubblicazione

Ogni articolo è consultabile nel sito internazionale digitando titolo e/o riferimenti rivista:
<https://pubmed.ncbi.nlm.nih.gov/>



Gli articoli scientifici sono aggiornati periodicamente (ogni 6 mesi) a partire dal più recente

Ulteriori pubblicazioni (Abstracts, Relazioni a Convegni Nazionali e Internazionali, Libri e Capitoli di Libro) sono a disposizione presso la Direzione della UOC



Eur J Cancer
2022 May;167:81-91.
doi: 10.1016/j.ejca.2022.03.005. Epub 2022 Mar 18.

Clinical profile and mortality of Sars-Cov-2 infection in cancer patients across two pandemic time periods (Feb 2020-Sep 2020; Sep 2020-May 2021) in the Veneto Oncology Network: The ROVID study

J Thorac Oncol
2022 Sep;17(9):1086-1097.
doi: 10.1016/j.jtho.2022.05.008. Epub 2022 Jun 1.

Addition of Bevacizumab to Erlotinib as First-Line Treatment of Patients With EGFR-Mutated Advanced Nonsquamous NSCLC: The BEVERLY Multicenter Randomized Phase 3 Trial

Target Oncol
2022 Jul;17(4):467-474.
doi: 10.1007/s11523-022-00892-z. Epub 2022 Jun 25.

Real-World Treatment with Nivolumab or Cabozantinib for Metastatic Renal Cell Carcinoma (mRCC) in the Veneto Region of Italy: Results of AMOUR Study

ESMO Open
2022 Apr;7(2):100431.
doi: 10.1016/j.esmoop.2022.100431. Epub 2022 Apr 8.

Real-world experience of abiraterone acetate plus prednisone in chemotherapy-naive patients with metastatic castration-resistant prostate cancer: long-term results of the prospective ABItude study

Eur J Cancer
2021 Dec;159:194-204.
doi: 10.1016/j.ejca.2021.09.005. Epub 2021 Nov 11.

Nasopharyngeal cancer in non-endemic areas: Impact of treatment intensity within a large retrospective multicentre cohort

Eur J Cancer
2021 Apr;147:120-127.
doi: 10.1016/j.ejca.2021.01.021. Epub 2021 Feb 3.

Epidemiology and clinical course of severe acute respiratory syndrome coronavirus 2 infection in cancer patients in the Veneto Oncology Network: The Rete Oncologica Veneta covid19 study

Eur J Cancer
2021 Sep;155:56-63.
doi: 10.1016/j.ejca.2021.06.016. Epub 2021 Aug 3.

Docetaxel and prednisone with or without enzalutamide as first-line treatment in patients with metastatic castration-resistant prostate cancer: CHEIRON, a randomised phase II trial

Target Oncol
2021 Nov;16(6):823-838.
doi: 10.1007/s11523-021-00831-4. Epub 2021 Oct 18.

Making Treatment-Free Remission (TFR) Easier in Chronic Myeloid Leukemia: Fact-Checking and Practical Management Tools

Front Oncol
2021 May 26;11:638689.
doi: 10.3389/fonc.2021.638689. eCollection 2021.

Perspectives and Emotional Experiences of Patients With Chronic Myeloid Leukemia During ENEST Path Clinical Trial and Treatment-Free Remission: Rationale and Protocol of the Italian Substudy

Front Cardiovasc Med
2020 Oct 26;7:598384.
doi: 10.3389/fcvm.2020.598384. eCollection 2020.

Development of Cancer in Patients With Heart Failure: How Systemic Inflammation Can Lay the Groundwork

Cancer Metastasis Rev
2020 Jun;39(2):535-552.
doi: 10.1007/s10555-020-09869-8.

How progressive cancer endangers the heart: an intriguing and underestimated problem

Breast Care (Basel)
2020 Feb;15(1):30-37.
doi: 10.1159/000495469. Epub 2019 Apr 17.

Is There Still a Role for Endocrine Therapy Alone in HR+/HER2- Advanced Breast Cancer Patients? Results from the Analysis of Two Data Sets of Patients Treated with High-Dose Fulvestrant as First-Line Therapy in the Real-World Setting: The EVA and GIM-13 AMBRA Studies

Cancer Res
2020 Feb 1;80(3):624-638.
doi: 10.1158/0008-5472.CAN-19-1840. Epub 2019 Nov 13.

Association of Genomic Domains in *BRCA1* and *BRCA2* with Prostate Cancer Risk and Aggressiveness

Pain Pract
2019 Mar 27.
doi: 10.1111/papr.12784. Online ahead of print.

Oral prolonged-release Oxycodone-Naloxone: analgesic response, safety profile, and factors influencing the response in advanced cancer patients

Eur J Cancer
2019 Jul;115:4-12.
doi: 10.1016/j.ejca.2019.03.022. Epub 2019 May 10.

Clinical outcomes and prognostic factors in recurrent and/or metastatic head and neck cancer patients treated with chemotherapy plus cetuximab as first-line therapy in a real-world setting

Front Psychol
2019 Feb 20;10:329.
doi: 10.3389/fpsyg.2019.00329. eCollection 2019.

Chronic Myeloid Leukemia Patient's Voice About the Experience of Treatment-Free Remission Failure: Results From the Italian Sub-Study of ENESTPath Exploring the Emotional Experience of Patients During Different Phases of a Clinical Trial

Breast Cancer Res Treat
2019 Apr;174(2):433-442.
doi: 10.1007/s10549-018-05070-2. Epub 2018 Dec 7.

Metformin plus chemotherapy versus chemotherapy alone in the first-line treatment of HER2-negative metastatic breast cancer. The MYME randomized, phase 2 clinical trial

Clin Transl Oncol
2019 Jun;21(6):790-795.
doi: 10.1007/s12094-018-1989-y. Epub 2018 Nov 17.

Metronomic oral vinorelbine for the treatment of advanced non-small cell lung cancer: a multicenter international retrospective analysis

Lung Cancer
2019 Jul;133:62-68.
doi: 10.1016/j.lungcan.2019.05.009. Epub 2019 May 9.

Effect on quality of life of cisplatin added to single-agent chemotherapy as first-line treatment for elderly patients with advanced non-small cell lung cancer: Joint analysis of MILES-3 and MILES-4 randomised phase 3 trials

N Engl J Med
2019 Dec 19;381(25):2391-2402.
doi: 10.1056/NEJMoa1910962. Epub 2019 Sep 28.

Niraparib in Patients with Newly Diagnosed Advanced Ovarian Cancer

Oncologist
2019 Jun;24(6):e318-e326.
doi: 10.1634/theoncologist.2018-0712. Epub 2019 Mar 7.

From Diagnostic-Therapeutic Pathways to Real-World Data: A Multicenter Prospective Study on Upfront Treatment for EGFR-Positive Non-Small Cell Lung Cancer (MOST Study)

Clin Breast Cancer
2019 Apr;19(2):137-145.e4.
doi: 10.1016/j.clbc.2018.11.009. Epub 2018 Nov 24.

A New Genetic Risk Score to Predict the Outcome of Locally Advanced or Metastatic Breast Cancer Patients Treated With First-Line Exemestane: Results From a Prospective Study

Lancet Oncol
2019 Sep;20(9):1273-1285.
doi: 10.1016/S1470-2045(19)30395-X. Epub 2019 Jul 22.

Adjuvant chemoradiotherapy versus radiotherapy alone in women with high-risk endometrial cancer (PORTEC-3): patterns of recurrence and post-hoc survival analysis of a randomised phase 3 trial

Br J Cancer
2019 Oct;121(9):744-750.
doi: 10.1038/s41416-019-0584-5. Epub 2019 Sep 20.

Multicenter, randomised, open-label, non-comparative phase 2 trial on the efficacy and safety of the combination of bevacizumab and trabectedin with or without carboplatin in women with partially platinum-sensitive recurrent ovarian cancer

Cancers (Basel) 2018 Jun 1;10(6):175.
doi: 10.3390/cancers10060175.

Factors Influencing the Clinical Presentation of Breakthrough Pain in Cancer Patients

J Clin Oncol
2018 Mar 1;36(7):689-696.
doi: 10.1200/JCO.2017.74.1652. Epub 2017 Nov 2.

Long-Term Results of the FOLL05 Trial Comparing R-CVP Versus R-CHOP Versus R-FM for the Initial Treatment of Patients With Advanced-Stage Symptomatic Follicular Lymphoma

BMC Cancer
2018 Jan 25;18(1):98.
doi: 10.1186/s12885-018-4001-x.

The PANDA study: a randomized phase II study of first-line FOLFOX plus panitumumab versus 5FU plus panitumumab in RAS and BRAF wild-type elderly metastatic colorectal cancer patients

Oncotarget
2018 Nov 30;9(94):36720-36721.

Correction 2: Everolimus (EVE) and exemestane (EXE) in patients with advanced breast cancer aged \geq 65 years: new lessons for clinical practice from the EVA study

J Clin Oncol
2018 Sep 1;36(25):2585-2592.
doi: 10.1200/JCO.2017.76.8390. Epub 2018 Jul 20.

Cisplatin-Based First-Line Treatment of Elderly Patients With Advanced Non-Small-Cell Lung Cancer: Joint Analysis of MILES-3 and MILES-4 Phase III Trials

Oncotarget
2018 Aug 7;9(61):31877-31887.
doi: 10.18632/oncotarget.25874.

Everolimus (EVE) and exemestane (EXE) in patients with advanced breast cancer aged \geq 65 years: new lessons for clinical practice from the EVA study

Lancet Oncol
2018 Mar;19(3):295-309.
doi: 10.1016/S1470-2045(18)30079-2. Epub 2018 Feb 12.

Adjuvant chemoradiotherapy versus radiotherapy alone for women with high-risk endometrial cancer (PORTEC-3): final results of an international, open-label, multicentre, randomised, phase 3 trial

Br J Cancer
2018 Aug;119(5):565-571.
doi: 10.1038/s41416-018-0190-y. Epub 2018 Jul 30.

A phase II randomised (calibrated design) study on the activity of the single-agent trabectedin in metastatic or locally relapsed uterine leiomyosarcoma

Oncotarget
2018 Oct 2;9(77):34639-34640.
doi: 10.18632/oncotarget.26213.

Correction: Everolimus (EVE) and exemestane (EXE) in patients with advanced breast cancer aged \geq 65 years: new lessons for clinical practice from the EVA study

Leuk Res 2018 Nov;74:75-79.
doi: 10.1016/j.leukres.2018.09.018. Epub 2018 Oct 2.

Safety and efficacy of switching from branded to generic imatinib in chronic phase chronic myeloid leukemia patients treated in Italy

Oncotarget
2018 Feb 12;9(18):14219-14227.
doi: 10.18632/oncotarget.24478. eCollection 2018 Mar 6.

Excellent outcomes of 2G-TKI therapy after imatinib failure in chronic phase CML patients

Leuk Res
2018 Nov;74:75-79.
doi: 10.1016/j.leukres.2018.09.018. Epub 2018 Oct 2.

Safety and efficacy of switching from branded to generic imatinib in chronic phase chronic myeloid leukemia patients treated in Italy

Lancet Haematol
2017 Mar;4(3):e137-e146.
doi: 10.1016/S2352-3026(17)30016-9. Epub 2017 Feb 17.

Lenalidomide maintenance in patients with relapsed diffuse large B-cell lymphoma who are not eligible for autologous stem cell transplantation: an open label, single-arm, multicentre phase 2 trial

Ann Oncol
2017 Sep 1;28(9):2206-2212.
doi: 10.1093/annonc/mdx299.

Induction TPF followed by concomitant treatment versus concomitant treatment alone in locally advanced head and neck cancer. A phase II-III trial

Adv Ther
2017 Jan;34(1):120-135.
Doi: 10.1007/s12325-016-0440-4. Epub 2016 Nov 21.

Breakthrough Cancer Pain: Preliminary Data of The Italian Oncologic Pain Multisetting Multicentric Survey (IOPS-MS)

Oncol Lett
2017 Jul;14(1):185-193.
doi: 10.3892/ol.2017.6152. Epub 2017 May 10.

Multicenter research into the quality of life of patients with advanced oropharyngeal carcinoma with long-term survival associated with human papilloma virus

Eur Arch Otorhinolaryngol
2017 Jun;274(6):2529-2536.
doi: 10.1007/s00405-017-4515-x. Epub 2017 Mar 10.

Follow-up of oral and oropharyngeal cancer using narrow-band imaging and high-definition television with rigid endoscope to obtain an early diagnosis of second primary tumors: a prospective study

Support Care Cancer
2017 Aug;25(8):2673-2674.
doi: 10.1007/s00520-017-3739-z.

Erratum to: Breakthrough pain and its treatment: critical review and recommendations of IOPS (Italian Oncologic Pain Survey) expert group

Clin Lung Cancer
2017 Jul;18(4):381-387.
doi: 10.1016/j.clcc.2016.10.002. Epub 2016 Nov 22.

Patients' Attitudes and Physicians' Perceptions Toward Maintenance Therapy for Advanced Non-Small-cell Lung Cancer: A Multicenter Italian Survey

Oncologist
2017 Jun;22(6):648-654.
doi: 10.1634/theoncologist.2016-0461. Epub 2017 Apr 21.

Everolimus Plus Exemestane in Advanced Breast Cancer: Safety Results of the BALLET Study on Patients Previously Treated Without and with Chemotherapy in the Metastatic Setting

Breast

2017 Oct;35:115-121.

doi: 10.1016/j.breast.2017.06.043. Epub 2017 Jul 13.

Efficacy and safety of Everolimus and Exemestane in hormone-receptor positive (HR+) human-epidermal-growth-factor negative (HER2-) advanced breast cancer patients: New insights beyond clinical trials. The EVA study

Adv Ther

2017 Jan;34(1):120-135.

doi: 10.1007/s12325-016-0440-4. Epub 2016 Nov 21.

Breakthrough Cancer Pain: Preliminary Data of The Italian Oncologic Pain Multisetting Multicentric Survey (IOPS-MS)

Oncologist

2017 Jun;22(6):648-654.

doi: 10.1634/theoncologist.2016-0461. Epub 2017 Apr 21.

Everolimus Plus Exemestane in Advanced Breast Cancer: Safety Results of the BALLET Study on Patients Previously Treated Without and with Chemotherapy in the Metastatic Setting

Support Care Cancer

2017 Jun;25(6):1743-1748.

doi: 10.1007/s00520-016-3547-x. Epub 2017 Jan 19.

Emesis and nausea related to single agent trabectedin in ovarian cancer patients: a sub-study of the MITO15 project

Haematologica

2017 May;102(5):e203-e206.

doi: 10.3324/haematol.2016.154211. Epub 2017 Jan 12.

Second-line rituximab, lenalidomide, and bendamustine in mantle cell lymphoma: a phase II clinical trial of the Fondazione Italiana Linfomi

Ann Oncol

2016 Jun;27(6):1107-15.

doi: 10.1093/annonc/mdw097. Epub 2016 Mar 2.

Are strong opioids equally effective and safe in the treatment of chronic cancer pain? A multicenter randomized phase IV 'real life' trial on the variability of response to opioids

Pharmacogenomics J
2016 Nov;16(6):525-529.
doi: 10.1038/tpj.2015.73. Epub 2015 Oct 27.

Predictive markers in elderly patients with estrogen receptor-positive breast cancer treated with aromatase inhibitors: an array-based pharmacogenetic study

Tumori
2016 Nov 11;102(Suppl. 2).
doi: 10.5301/tj.5000498.

Unusual cardiac metastasis of uterine leiomyosarcoma: case report and literature review

Cancer Metastasis Rev
2016 Sep;35(3):413-26.
doi: 10.1007/s10555-016-9625-1.

Targeting cellular and molecular drivers of head and neck squamous cell carcinoma: current options and emerging perspectives

Eur Urol
2015 Jul;68(1):147-53.
doi: 10.1016/j.eururo.2014.10.014. Epub 2014 Oct 25.

Clinical Outcomes of Castration-resistant Prostate Cancer Treatments Administered as Third or Fourth Line Following Failure of Docetaxel and Other Second-line Treatment: Results of an Italian Multicentre Study

Crit Rev Oncol Hematol
2015 Apr;94(1):98-104.
doi: 10.1016/j.critrevonc.2014.10.013. Epub 2014 Nov 7.

Rare uterine cancer: carcinosarcomas. Review from histology to treatment

Clin Lymphoma Myeloma Leuk
2015 Oct;15(10):592-8.
doi: 10.1016/j.clml.2015.05.003. Epub 2015 Jun 6.

Cytogenetic Impact on Lenalidomide Treatment in Relapsed/Refractory Multiple Myeloma: A Real-Life Evaluation

Anticancer Res
2015 Apr;35(4):2183-9.

Epirubicin plus paclitaxel regimen as second-line treatment of patients with small-cell lung cancer

Expert Opin Pharmacother
2013 Oct;14(15):2011-7.
doi: 10.1517/14656566.2013.829041.

Is pain reporting to health care professionals age-related? A cross sectional multicenter study in a hospital setting

BMC Res Notes
2012 Sep 3;5:482.
doi: 10.1186/1756-0500-5-482.

Re-challenge with pemetrexed in advanced mesothelioma: a multi-institutional experience

J Thorac Oncol
2012 Jan;7(1):233-42.
doi: 10.1097/JTO.0b013e318233d6c2.

Phase I-II trial of gemcitabine-based first-line chemotherapies for small cell lung cancer in elderly patients with performance status 0-2: the G-STEP trial

Eur J Gynaecol Oncol
2011;32(3):328-30.

Laparoscopic total fallopian tube removal at the time of bilateral salpingo-oophorectomy in BRCA2 positive women

J Clin Oncol
2010 Feb 10;28(5):866-71.
doi: 10.1200/JCO.2009.23.6125. Epub 2009 Dec 28.

Genotype-driven phase I study of irinotecan administered in combination with fluorouracil/leucovorin in patients with metastatic colorectal cancer

Eur J Gynaecol Oncol
2010;31(6):658-61.

Overall survival in BRCA-associated ovarian cancer: case-control study of an Italian series

Br J Cancer
2009 May 19;100(10):1566-74.
doi: 10.1038/sj.bjc.6605053. Epub 2009 Apr 28.

Pattern and quality of care of cancer pain management. Results from the Cancer Pain Outcome Research Study Group

Assist Infirm Ric
2009;28(2):73-81.

[The ECAD_O project: clinical caring epidemiology of pain in the hospital: the nurses' contribution]

Oncologist
2009 May;14(5):497-510.
doi: 10.1634/theoncologist.2008-0260. Epub 2009 May 7.

Impact of third-generation drugs on the activity of first-line chemotherapy in advanced non-small cell lung cancer: a meta-analytical approach

Clin J Pain
2009 Oct;25(8):671-82.
doi: 10.1097/AJP.0b013e3181a38f9d.

Effects of transdermal buprenorphine on patients-reported outcomes in cancer patients: results from the Cancer Pain Outcome Research (CPOR) Study Group

Biol Blood Marrow Transplant
2009 Mar;15(3):326-35.
doi: 10.1016/j.bbmt.2008.11.036.

Reduced-intensity allogeneic hematopoietic stem cell transplantation in metastatic colorectal cancer as a novel adoptive cell therapy approach. The European group for blood and marrow transplantation experience

Lung Cancer
2008 Jul;61(1):67-72.
doi: 10.1016/j.lungcan.2007.12.002.

The MILES-2G phase 2 study of single-agent gemcitabine with prolonged constant infusion in advanced non-small cell lung cancer elderly patients

Clin Drug Investig
2008;28(7):399-407.
doi: 10.2165/00044011-200828070-00001.

Oxycodone controlled-release as first-choice therapy for moderate-to-severe cancer pain in Italian patients: results of an open-label, multicentre, observational study

J Card Fail
2007 Nov;13(9):701-8.
doi: 10.1016/j.cardfail.2007.06.722.

Effects of exercise training on endothelial progenitor cells in patients with chronic heart failure

Lancet Oncol
2007 Jun;8(6):500-12.
doi: 10.1016/S1470-2045(07)70146-8.

Factorial phase III randomised trial of rofecoxib and prolonged constant infusion of gemcitabine in advanced non-small-cell lung cancer: the GEMcitabine-COXib in NSCLC (GECO) study

Aging Clin Exp Res
2007 Apr;19(2):91-6.
doi: 10.1007/BF03324673.

Amino-bisphosphonates decrease hTERT gene expression in breast cancer in vitro

Breast Cancer Res Treat
2007 Mar;101(3):279-84.
doi: 10.1007/s10549-006-9300-2. Epub 2006 Jul 12.

Correlation between genetic and biological aspects in primary non-metastatic breast cancers and corresponding synchronous axillary lymph node metastasis

Am Heart J
2006 Mar;151(3):674-80.
doi: 10.1016/j.ahj.2005.04.033.

Direct intramyocardial percutaneous delivery of autologous bone marrow in patients with refractory myocardial angina

BMC Cancer
2006 Mar 3;6:49.
doi: 10.1186/1471-2407-6-49.

The effect of bisphosphonates on gene expression: GAPDH as a housekeeping or a new target gene?

BMC Cancer
2006 Feb 27;6:44.
doi: 10.1186/1471-2407-6-44.

Tolerability of intensified intravenous interferon alfa-2b versus the ECOG 1684 schedule as adjuvant therapy for stage III melanoma: a randomized phase III Italian Melanoma Inter-group trial (IMI - Mel.A.) [ISRCTN75125874]

Diagn Mol Pathol
2006 Dec;15(4):206-15.
doi: 10.1097/01.pdm.0000213469.85301.d6.

Pattern and distribution of immunoglobulin VH gene usage in a cohort of B-CLL patients from a Northeastern region of Italy

Acta Otorhinolaryngol Ital
2005 Jun;25(3):195-9.

Maxillo-ethmoidal chloroma in acute myeloid leukaemia: case report

Mol Cell Endocrinol
2005 Aug 30;240(1-2):23-31.
doi: 10.1016/j.mce.2005.03.018.

Bisphosphonates decrease telomerase activity and hTERT expression in MCF-7 breast cancer cells

Lancet Oncol
2005 Sep;6(9):669-77.
doi: 10.1016/S1470-2045(05)70255-2.

Chemotherapy-induced neutropenia and treatment efficacy in advanced non-small-cell lung cancer: a pooled analysis of three randomised trials

J Clin Oncol
2005 Oct 1;23(28):6865-72.
doi: 10.1200/JCO.2005.02.527.

Pretreatment quality of life and functional status assessment significantly predict survival of elderly patients with advanced non-small-cell lung cancer receiving chemotherapy: a prognostic analysis of the multicenter Italian lung cancer in the elderly study

Int J Biol Markers
2005;20(1):28-33.
doi: 10.5301/jbm.2008.5136.

A new nested primer pair improves the specificity of CK-19 mRNA detection by RT-PCR in occult breast cancer cells

Br J Cancer
2004 Jun 14;90(12):2288-96.
doi: 10.1038/sj.bjc.6601810.

Prevalence and management of pain in Italian patients with advanced non-small-cell lung cancer

Br J Cancer
2003 Sep 15;89(6):1013-21.
doi: 10.1038/sj.bjc.6601236.

Supportive care in patients with advanced non-small-cell lung cancer

Stem Cells
2003;21(1):33-40.
doi: 10.1634/stemcells.21-1-33.

The impact of progenitor enrichment, serum, and cytokines on the ex vivo expansion of mobilized peripheral blood stem cells: a controlled trial

J Natl Cancer Inst
2003 Mar 5;95(5):362-72.
doi: 10.1093/jnci/95.5.362.

Chemotherapy for elderly patients with advanced non-small-cell lung cancer: the Multicenter Italian Lung Cancer in the Elderly Study (MILES) phase III randomized trial

Histochem J
2002;34(8-9):403-10.
doi: 10.1023/a:1023635521069.

Human fibroblasts from normal and malignant breast tissue grown in vitro show a distinct senescence profile and telomerase activity

Histochem J
2001;33(9-10):499-509.
doi: 10.1023/a:1014927305775.

Conditioned medium from MCF-7 cell line induces myofibroblast differentiation, decreased cell proliferation, and increased apoptosis in cultured normal fibroblasts but not in fibroblasts from malignant breast tissue

Oncology
2000 Aug;59(2):100-4.
doi: 10.1159/000012144.

Hypnosis in the treatment of anticipatory nausea and vomiting in patients receiving cancer chemotherapy

J Cancer Res Clin Oncol
1998;124(2):93-105.
doi: 10.1007/s004320050140.

Differentiation, proliferation and apoptosis levels in human leiomyoma and leiomyosarcoma

J Muscle Res Cell Motil
1997 Aug;18(4):413-27.
doi: 10.1023/a:1018642713934.

Oestrogen-dependent expression of the SM2 smooth muscle-type myosin isoform in rabbit myometrium

Eur Urol
1996;29(3):312-6; discussion 317.
doi: 10.1159/000473767.

Carboplatin monochemotherapy in elderly patients with nonoperable transitional cell carcinoma of the bladder: a two-stage, phase II study

Eur J Cancer
1996 Aug;32A(9):1612-5.
doi: 10.1016/0959-8049(96)00134-7.

The impact of a psychological intervention on quality of life in non-metastatic breast cancer

Cell Tissue Res
1996 Jan;283(1):7-18.
doi: 10.1007/s004410050507.

Expression of non-muscle myosin isoforms in rabbit myometrium is estrogen-dependent

Oncology
1993;50(1):1-4.
Doi: 10.1159/000227137.

The combination of mitomycin, vinblastine and cisplatin is active in the palliation of stage IIIB-IV non-small-cell lung cancer
